

# The Journal of Phytopharmacology

(Pharmacognosy and phytomedicine Research)

## Research Article

ISSN 2320-480X  
 JPHYTO 2018; 7(1): 13-18  
 January- February  
 Received: 24-04-2017  
 Accepted: 10-12-2017  
 © 2018, All rights reserved

### Kajal Kumari

School of Bioengineering and Food  
 Technology, Shoolini University, Solan,  
 Himachal Pradesh, India

### Somesh Sharma

School of Bioengineering and Food  
 Technology, Shoolini University, Solan,  
 Himachal Pradesh, India

### V.K. Joshi

School of Bioengineering and Food  
 Technology, Shoolini University, Solan,  
 Himachal Pradesh, India

### Surabhi Sharma

College of Agriculture Sciences,  
 Chandigarh University, Ropar, Punjab,  
 India

### Correspondence:

#### Somesh Sharma

School of Bioengineering and Food  
 Technology, Shoolini University, Solan,  
 Himachal Pradesh, India

Email:

someshsharma[at]shooliniuniversity.com

## Adding value to wild Himalayan fig (*Ficus palmata*): Composition, functional and sensory characteristics of jam

Kajal Kumari, Somesh Sharma, V.K. Joshi, Surabhi Sharma

### ABSTRACT

Wild Himalayan fig is an underutilized wild fruit with various nutritional components. The fruit was evaluated for different parameters like crude fibre value (18.90 - 16.38 %), total phenols (72.6 – 65.4 mg/100g) and the energy value (99.84 Kcal). The shelf life of this fruit was enhanced by converting it into value added product like jam. To standardize the technology for jam preparation, different treatments were designed by varying the concentration of pectin and citric acid. Among these treatments, T<sub>3</sub>C<sub>1</sub> containing 0.7 percent pectin and 0.3 percent citric acid was found best with higher titratable acidity (12.02%) as citric acid, ascorbic acid content (1.24mg/100g), total sugar (66.42 %) and reducing sugar (58.85%), total phenol (39.8 mg/100g), anthocyanin 17.05 mg/100 g). Further, on the basis of sensory evaluation, the standardized product had high overall acceptability along with higher score for colour, flavor etc. The FTIR analysis of prepared jam and fig pulp clearly revealed that there was no alteration in physicochemical properties of fig pulp after its processing. Hence, the results from the present investigation clearly revealed that development of jam from wild fig can also be one of the alternative for the utilization of this nutritional rich underutilized crop for commercial purposes.

**Keywords:** Jam, *Ficus palmata*, Wild Himalayan Fig.

### INTRODUCTION

*Ficus palmata* belongs to family Rosaceae is one of the important wild fruits found growing in western Himalayan regions of India. The tree is deciduous in nature and found in the forests, around the villages in wastes lands or on the bunds of farmer fields<sup>[1, 2]</sup>. The wild Himalayan fig is commonly found in states of Uttarakhand, Punjab, Kashmir and areas of western Himalayan region<sup>[1]</sup>. Fresh fig fruits weight, length and breadth ranged between 11.6-36.8 g, 16.7-28.6 cm and 17.3-31.0 cm, respectively<sup>[3]</sup>. Figs are an excellent source of phytosterols<sup>[4]</sup>, crude fibre, minerals<sup>[5]</sup>, carotenoid<sup>[6]</sup>, anthocyanins<sup>[7-9]</sup> and polyphenol<sup>[8, 10]</sup>. The fig is used in various diseases e.g. gastrointestinal, hypoglycemic, anti-tumour, anti-ulcer, anti-diabetic, lipid lowering and antifungal activities<sup>[11]</sup>. However, there is no documented data available on the production of this fruit. Till date, the fruits are underutilized and farmers are even cutting these fruit trees as they are unawareness about the nutritional quality characteristics of the fruit. Furthermore, the fruit is edible along with the seeds and has high medicinal value along with sensory scores<sup>[2, 12]</sup>. However, presently the green fruits of the wild Himalayan fig are utilized as cooked vegetable by the rural people of these regions<sup>[1]</sup>. It can also be used for making various products such as squash, jam and jelly<sup>[12, 13]</sup>. The fruits are safe for human consumption and may possess high value components such as sugars, phenols, antioxidants and fibre that may be used in development of value added products with high nutritive value<sup>[1]</sup>. Saklani and Chandra (2011b) reported that bark, root, leaves, fruit and latex of this plant are used for the various illnesses treatment<sup>[14]</sup>. Solomon *et al.* (2006) reported that fig fruit skin was richest source of anthocyanin and polyphenol in comparison to other parts of fig fruit<sup>[7]</sup>. Glimpses of present research conducted into various fig products have not been a true reflection of the importance and potential of this crop.

Many processes designed to preserve food will involve a number of food preservation methods. Among these, Jam is one of the most common techniques. In India, since the fruit is underutilized no work has been done on development and evaluation of wild Himalayan fig jam for its commercialization. To enhance the value of *Ficus palmata*, one should begin with a survey of the physicochemical

characteristics of the raw material from the local material. Subsequently, laboratory assays may be conducted in order to prepare and evaluate *Ficus palmata* products with high added value such as fig jam. Further, the knowledge of physico-chemical characteristics of the final products is also essential for conception of equipment size and process. Hence, this paper reports on the value addition of fig from the underutilized species *Ficus palmata* through characterization of the raw material. Further, the development and evaluation of wild Himalayan fig would provide sustainable means to the local people of these regions. This would also be helpful for farmers and industries to utilize and design new food products, machinery and quality control for commercialization of such products.

## MATERIALS AND METHODS

### Collection of *Ficus palmata* fruits

This study was conducted on wild Himalayan fig (*Ficus palmata*) which is one of the underutilized fruit in mid-Hill region of Himachal Pradesh, India. Fully ripened fruits of wild Himalayan fig were collected from different districts (Hamirpur, Bilaspur, Kangra, Solan). The fruits were washed and sorted for the presence of any defects and packed in polyethylene bags and kept at 4°C until analysed/utilized.

### Preparation of pulp

The fig pulp was prepared by boiling the fruits with water (1:1 ratio) for 15 mins.

### Preparation of Wild Himalayan fig jam

Preserving fruit by turning it into jam, for example, involves boiling (to reduce the fruit's moisture content and to kill bacteria, yeasts, etc.), addition of sugar (to prevent their re-growth) and sealing within an airtight jar (to prevent recontamination). Different treatments of jam were prepared using varying concentration of citric acid (0.3 to 0.7 %) and pectin (0.3 to 0.7 %). The jam was prepared by adding sucrose to raise the final TSS to 68° Brix in an open pan, with continuous stirring. Finally, when the TSS reached to final concentration, the jam was poured in pre sterilized glass jars. The jars were cooled to room temperature and stored at ambient conditions (RT 25-27°C) till further evaluation of various physico-chemical, storage and sensory characteristics.

### Physico-chemical analysis of Wild Himalayan fig and jam

Total soluble solids content (Brix) was determined using an Abbe refractometer at 25°C [15]. Titratable acidity in terms of percent citric acid was determined according to the AOAC methods [16]. Total and reducing sugar content was determined as per the method described by Rangana [17]. Ascorbic acid content was measured as mg/100 g using 2,6-Dichlorophenol-Indophenol visual titration [17]. Total phenolic content was determined at 725 nm using Folin-Ciocalteu reagent method [18] and expressed as gallic acid equivalents (GAE mg/100 g of sample, fresh weight). Crude fibre content was measured according to the method given by Rangana [17]. Anthocyanin content was determined as per the method described by Thimmaiah [19]. Further, antioxidant activity was measured as % DPPH scavenging activity [20]. The water activity of Wild Himalayan Fig jam was measured at 25 °C in Rotronic HW4 apparatus (Rotronic instrument corp, New York). Further, the FTIR analysis of the prepared treatments was conducted in Agilent instrument.

### Sensory evaluation

The sensory evaluation of jam was conducted by semi-trained panel of 10 judges. Each judge was given a set of treatments separately in isolated booths and provided with a glass of fresh water to rinse their mouth before tasting the next sample. Each sample was evaluated for various quality attributes viz., colour, flavor, consistency/texture, taste/acid sugar blend and overall acceptability. Judges were asked to rate the samples on a prescribed sensory evaluation performa of composite scoring.

### Statistical analysis

Analytical values were determined, using three independent determinations. Values of different parameters were expressed as the mean  $\pm$  standard deviation ( $X \pm SD$ ). The data of quantitative estimation of various physico-chemical characteristics of different jam treatments was analyzed by Completely Randomized Design (CRD) as described by O'Mahony (1985).

### Cost of production

The economics of production (per unit) of developed jam at the laboratory scale was worked out based on the cost of all ingredients, 10 per cent overhead charges, 20 per cent processing charges (cost of making puree/pulp/slurry) and 20 per cent profit margin.

## RESULTS AND DISCUSSION

### Compositional and functional characteristics of Wild Himalayan Fig

The proximate composition of wild Himalayan Fig (ripened fruit) collected from different districts of Himachal Pradesh is presented in Table 1. It is clearly seen from the results that wild fig is a good source of crude fiber (18.90-18.03 %) with relatively low content of ascorbic acid (3.8 - 3.09 mg/100 g). The total soluble solid content of the fruit from different districts ranged between 12.8 to 11.2°B with highest in samples collected from District Hamirpur and minimum in fruits from Bilaspur. Since, total soluble solids are one of the important quality attribute for the raw consumption and processing of fruits. Hence, the fruits can be effectively used for the development of value added products from wild fig. Among the different districts, the titratable acidity content was recorded between 0.44 - 0.38 per cent as citric acid. The maximum value for titratable acidity was recorded in samples collected from Solan and minimum in fruits from Hamirpur. Further, wild figs also contain total phenols (72.6 – 65.4 mg/100 g) although the quantity is less in comparison to cultivated fig varieties. Indeed, they were rich in sugars and fiber deserving to be used for value-added product formulation. The perusal of the data further revealed that the protein content and reducing sugar of fruits collected from different locations ranged between 1.35-1.82 per cent and 4.17 – 3.06 mg/100g with 0.20-0.17 percent pectin. The results on energy value (99.84 – 94.26 Kcal) of fruit also signify that wild fig fruit is a good source of energy. There were significant differences among the fruits collected from different locations of the Himachal Pradesh for presence of antioxidants (Table 2). The highest antioxidant activity (78.25%) by DPPH method was recorded in fruits

collected from Distt Hamirpur. Further, the results pertaining to FRAP and ABTS activity also support the presence of higher antioxidants in fruits collected from Hamirpur District of Himachal Pradesh. Furthermore, the similar trend was also recorded in ABTS activity of Wild Himalayan fig collected from different locations.

**Chemical Characteristics of wild Himalayan fig Jam**

The perusal of data in Table 3 revealed that among the various treatments, T<sub>3</sub>C<sub>1</sub> (0.7%+0.3%) was found best having higher titratable acidity (0.56 asper cent citric acid), ascorbic acid content (1.24 mg/100g), total sugar (66.42 %) and reducing sugar (58.85 %) respectively.

Further, Table 4 revealed that there were significant differences among the treatments for total phenols which ranges from 39.8 to 37.2 mg/100g. The highest content of phenols was recorded in treatment T<sub>3</sub>C<sub>1</sub> whereas, non-significant differences were observed in anthocyanins and % DPPH scavenging activity respectively. Further, the setting of all the jams was recorded at 68-71 °B. The increase in reducing sugars content in jam compared with the raw material could be explained by a partial hydrolysis of sucrose (initial and added) during cooking (Pearson, 1976). The reducing sugars, which have more affinity for water than sucrose, could contribute to the reduction of the jam crystallization phenomenon [21]. The cost of production of fig jam was worked out to be Rs. 28 per 250 gjar (Table 5) which was found to economical when compared with

other jams in the market.

**FTIR analysis of prepared jam and pulp**

The results pertaining to the FFTIR analysis have been shown in Table 6. The grouping present in the pulp having same peak at particular wavelength as in prepared jam. The various functional groups present (Fig 1 and 2) in both fig pulp and jam were (-C=C-O-C) (Ether), (-OH) (Hydroxyl), (-O-NO<sub>2</sub>) (3° Amide), (-C-OH) (Phenol) and (-C-OH) (1° and 2° Alcohol).

The presence of these functional groups represents the molecular composition of both pulp and jam which further derives the conclusion that there is no interaction in characteristic functional groups of pulp with the ingredients of jam. Therefore, no alteration in physiochemical properties were observed.

**Sensory characteristics of wild Himalayan fig jam**

Fig jam was characterized by a better colour, texture and flavour. Different treatments of fig jam were presented to panel of judges for sensory evaluation on composite scoring performa and the results (Table 7 and Fig 3) showed the highest overall acceptability scores of 81.25 out of 100 were awarded to treatment T<sub>3</sub>C<sub>1</sub> along with highest scores for other characteristics also such as colour, flavor, sugar acid blend etc.

**Table 1:** Proximate Composition of fresh Wild Himalayan fig from various Districts of Himachal Pradesh

Parameters	Quantity				CD <sub>(0.05)</sub>
	Hamirpur	Bilaspur	Kangra	Solan	
Moisture (%)	76.8	74.2	79.5	76.4	1.18
TSS (°B)	12.8	11.2	12.1	11.9	0.17
Titratable acidity (%)	0.380	0.440	0.421	0.445	0.01
Protein (%)	1.35	1.41	1.68	1.82	0.17
Total sugar (%)	7.01	6.56	6.12	6.41	0.12
Reducing sugar (%)	4.17	4.16	4.06	3.06	0.15
Total phenols (mg/100g)	72.6	65.4	68.9	70.6	5.42
Ascorbic acid (mg/100g)	3.67	3.2	3.09	3.8	0.19
Pectin (%)	0.20	0.17	0.19	0.18	NS
Crude fibre (%)	18.90	18.03	18.78	18.61	NS
Crude fat (%)	1.07	1.15	0.69	0.78	0.14
Ash (%)	0.92	0.84	0.72	0.98	NS
Anthocyanins (A <sub>525</sub> )	0.181	0.179	0.176	0.180	NS
Carbohydrate (mg/100g)	19.84	18.40	19.11	18.32	1.13
Energy (Kcal)	99.84	95.6	98.10	94.26	1.16

All values given are means of three determinations. Means in line with different letters are significantly different (P < 0.05).

**Table 2:** Antioxidant activity of fresh Wild Himalayan fig from different locations of Himachal Pradesh

Antioxidant Tests	Wild Himalayan fig				Mean	CD <sub>(0.05)</sub>
	Hamirpur	Bilaspur	Solan	Kangra		
DPPH (%)	78.25	76.51	76.87	78.03	77.41	1.13
FRAP [µmol Fe(II)/g extract]	987	856	896	971	927.5	2.72
ABTS (%)	92.05	89.01	89.23	91.04	90.33	0.179

All values given are means of three determinations. Means in line with different letters are significantly different (P < 0.05).

**Table 3:** Effect of different concentration of Pectin: Citric acid on Titratable acidity, ascorbic acid, total sugar, reducing sugar and setting of jam.

Treatments	Titratable Acidity (%)	Ascorbic Acid (mg/100g)	Total Sugar (%)	Reducing Sugar (%)
T <sub>1</sub> C <sub>1</sub> (0.3%+0.3%)	0.53±0.02 <sup>b</sup>	1.56±0.01 <sup>c</sup>	65.82±0.04 <sup>a</sup>	58.03±0.03 <sup>b</sup>
T <sub>1</sub> C <sub>2</sub> (0.3%+0.5%)	0.57±0.02 <sup>c</sup>	1.87±0.03 <sup>d</sup>	65.50±0.03 <sup>a</sup>	58.27±0.10 <sup>b</sup>
T <sub>1</sub> C <sub>3</sub> (0.3%+0.7%)	0.61±0.01 <sup>d</sup>	0.95±0.02 <sup>a</sup>	66.01±0.04	58.05±0.06 <sup>b</sup>
T <sub>2</sub> C <sub>1</sub> (0.5%+0.3%)	0.56±0.01 <sup>c</sup>	1.23±0.02 <sup>b</sup>	65.19±0.02 <sup>a</sup>	57.41±0.14 <sup>a</sup>
T <sub>2</sub> C <sub>2</sub> (0.5%+0.5%)	0.52±0.02 <sup>a</sup>	2.01±0.02 <sup>e</sup>	65.56±0.03 <sup>a</sup>	57.04±0.05 <sup>a</sup>
T <sub>2</sub> C <sub>3</sub> (0.5%+0.7%)	0.68±0.03 <sup>e</sup>	1.92±0.03 <sup>e</sup>	66.71±0.04 <sup>b</sup>	58.24±0.05 <sup>b</sup>
T <sub>3</sub> C <sub>1</sub> (0.7%+0.3%)	0.56±0.02 <sup>c</sup>	1.24±0.04 <sup>b</sup>	66.42±0.06 <sup>b</sup>	58.85±0.04 <sup>b</sup>
T <sub>3</sub> C <sub>2</sub> (0.7%+0.5%)	0.52±0.01 <sup>a</sup>	1.97±0.04 <sup>e</sup>	65.48±0.05 <sup>a</sup>	57.96±0.06 <sup>a</sup>
T <sub>3</sub> C <sub>3</sub> (0.7%+0.7%)	0.62±0.02 <sup>d</sup>	1.84±0.02 <sup>d</sup>	65.71±0.04	58.01±0.04
<b>Mean±SD</b>	<b>0.57±0.08</b>	<b>0.9±0.15</b>	<b>36.90±0.37</b>	<b>28.12±0.87</b>

Data are presented as mean±SD (n=3)

**Table 4:** Effect of different concentration of Pectin: Citric acid on Total Phenol, anthocyanin and % DPPH Scavenging activity of jam.

Treatments	Total phenol (mg/100g)	Anthocyanin (A <sub>525</sub> )	% DPPH Scavenging activity	Jam setting (°Brix)
T <sub>1</sub> C <sub>1</sub> (0.3%+0.3%)	37.2±0.10 <sup>a</sup>	0.066±0.11 <sup>c</sup>	27.51±0.18 <sup>c</sup>	71±0.03 <sup>c</sup>
T <sub>1</sub> C <sub>2</sub> (0.3%+0.5%)	38.3±0.04 <sup>b</sup>	0.062±0.19 <sup>b</sup>	26.82±0.11	71±0.04 <sup>c</sup>
T <sub>1</sub> C <sub>3</sub> (0.3%+0.7%)	38.7±0.02 <sup>b</sup>	0.067±0.05 <sup>c</sup>	26.52±0.05 <sup>b</sup>	71±0.03 <sup>c</sup>
T <sub>2</sub> C <sub>1</sub> (0.5%+0.3%)	37.4±0.01 <sup>a</sup>	0.068±0.04	27.63±0.05 <sup>c</sup>	69±0.01 <sup>b</sup>
T <sub>2</sub> C <sub>2</sub> (0.5%+0.5%)	39.2±0.12 <sup>c</sup>	0.062±0.06 <sup>b</sup>	22.96±0.14 <sup>a</sup>	69±0.03 <sup>b</sup>
T <sub>2</sub> C <sub>3</sub> (0.5%+0.7%)	37.5±0.6 <sup>a</sup>	0.059±0.01 <sup>a</sup>	25.05±0.17 <sup>b</sup>	69±0.01 <sup>b</sup>
T <sub>3</sub> C <sub>1</sub> (0.7%+0.3%)	39.8±0.4 <sup>c</sup>	0.067±0.08 <sup>c</sup>	27.53±0.15 <sup>c</sup>	68±0.03 <sup>a</sup>
T <sub>3</sub> C <sub>2</sub> (0.7%+0.5%)	38.2±0.9 <sup>b</sup>	0.064±0.15 <sup>b</sup>	23.89±0.18 <sup>a</sup>	68±0.04 <sup>a</sup>
T <sub>3</sub> C <sub>3</sub> (0.7%+0.7%)	39.1±0.01 <sup>c</sup>	0.063±0.06 <sup>b</sup>	26.71±0.17 <sup>b</sup>	68±0.01 <sup>a</sup>
<b>Mean±SD</b>	<b>38.4±2.94</b>	<b>0.064±0.003</b>	<b>26.06±1.70</b>	<b>69±0.03</b>

Data are presented as mean±SD (n=3)

**Table 5:** Cost of production of Jam:

Particulars	Quantity	Rate (Rs)	Amount (Rs)	
			(Glass jar)	(Plastic jar)
Fruit	500g	30/kg	15.0	15.0
Sugar	498.4g	40/kg	19.93	19.93
Sodium benzoate	0.8g	480/kg	0.38	0.38
Citric acid	3.0g	450/kg	1.35	1.35
Cardamom flavor	12 ml	15/25 ml	7.2	7.2
Pectin	5g	3500/500 g	35.0	35.0
Glass jar 250 g	1	9/jar	9	-
Plastic jar 250 g	1	15/jar	-	15
<b>Total cost</b>			87.86	93.86
<b>Overhead cost @10 %</b>			8.78	9.38
<b>Total cost</b>			96.64	103.24
<b>Profit @ 20%</b>			19.32	20.64
<b>Net cost (cost per 250 g jar)</b>			28.75	30.97


**Table 6:** Comparison of functional groups present in Fig pulp and jam

Sample	-C=C-O-C (Ether) Stretching	-OH (Hydroxyl) stretching	-O-NO <sub>2</sub> (3° Amide) Stretching	-C-OH (Phenol) Stretching	-C-OH (1° and 2° Alcohol) Stretching
Fig pulp	1030	3223	1640	1395	1298
Fig jam	1045	3231	1644	1339	1268


\*Expressed as wavenumber

**Table 7:** Sensory properties of standardized Fig jam:


Quality attributes	Score	Samples								
		T <sub>1</sub> C <sub>1</sub>	T <sub>1</sub> C <sub>2</sub>	T <sub>1</sub> C <sub>3</sub>	T <sub>2</sub> C <sub>1</sub>	T <sub>2</sub> C <sub>2</sub>	T <sub>2</sub> C <sub>3</sub>	T <sub>3</sub> C <sub>1</sub>	T <sub>3</sub> C <sub>2</sub>	T <sub>3</sub> C <sub>3</sub>
<b>Colour</b>	<b>20</b>	14.0	14.25	16.0	10.25	13.25	15.5	17.0	11.5	12.75
<b>Texture</b>	<b>20</b>	15.75	13.75	16.5	11.75	12.25	15.25	16.0	13.75	13.5
<b>Flavour</b>	<b>20</b>	14.5	14.15	14.25	12.75	11.5	15.75	16.75	13.0	14.5
<b>Sugar- acid blend</b>	<b>20</b>	14.0	14.75	18.0	12.0	11.25	15.5	15.75	11.25	13.0
<b>Absence of defects</b>	<b>20</b>	17.5	15.75	14.25	13.5	13.75	14.5	15.75	15.25	14.25
<b>Total score</b>	<b>100</b>	75.75	72.75	78.75	60.25	62.0	76.75	81.25	64.75	68.0



**Figure 1:** Graphical representation of FTIR analysis of Fig pulp



**Figure 2:** Graphical representation of FTIR analysis of fig jam



**Figure 3:** Overall acceptability scores of Wild Himalayan fig jam

**CONCLUSION**

It can be concluded from the study that wild Himalayan fig is one of the nutritious underutilized fruit crop grown in the Himachal region of western Himalayas. The fruit has higher fibre content with optimum amount of other nutrients such as sugars, phenols, antioxidants, anthocyanins etc. and low ascorbic acid. The utilization of these wild

fruits for jam making can be one of the alternative for adding value to the crop and as an income source to the farmers. Further, the knowledge of the physico-chemical and sensory properties of this product will also encourage the farmers to develop the products on an industrial scale. The use of fig jam may also be attractive to consumers as a positive alternative to conventional fruit jams available in the market.

**REFERENCES**

1. Parmar C, Kaushal MK. *Ficus palmate* in Wild Fruits. Kalyani Publishers, New Delhi, India. 1982; 31-34.
2. Chopra RN, Nayar SL, Chopra IC. Glossary of Indian Medicinal Plants (Including the Supplement). Council of Scientific and Industrial Research, New Delhi, 1986.
3. Sharma SK, Badiyala SD. Variability studies in common fig in Hamirpur district of Himachal Pradesh. Indian Journal of Horticulture. 2006; 63(2):159-161.
4. Jeong WS, Lachance PA. Phytosterols and fatty acids in fig (*Ficus carica* var. Mission) fruit and tree components. Food and Chemical Toxicology. 2001; 66:278-281.
5. USDA Nutritive Value of Foods., U.S. Department of Agriculture, Agricultural Research Service, Nutrient Data Laboratory, Beltsville, Maryland. Available at: [http://www.nal.usda.gov/fnic/foodcomp/Data/HG72/hg72\\_2002.pdf](http://www.nal.usda.gov/fnic/foodcomp/Data/HG72/hg72_2002.pdf) (Accessed 3 December 2008).
6. Su Q, Rowley KG, Itsiopoulos C, O'Dea K. Identification and quantitation of major carotenoids in selected components of the Mediterranean diet: green leafy vegetables, figs and olive oil. Eur J Clin Nutr. 2002; 56(11):1149-54.
7. Solomon A, Golubowicz S, Yablowicz Z. Antioxidant activities and anthocyanin content of fresh fruits of common fig (*Ficus carica* L.). Journal of Agricultural and Food Chemistry. 2006; 54:7717-7723.
8. Del Caro A, Piga A. Polyphenol composition of peel and pulp of two Italian fresh fig fruits cultivars (*Ficus carica* L.). European Food Research and Technology. 2008; 226:715-719.
9. Duenas M, Perez-Alonso JJ, Santos-Buelga C, Escribano-Bailon T. Anthocyanin composition in fig (*Ficus carica* L.). Journal of Food Composition and Analysis. 2008; 21:107-115.
10. Veberic R, Colaric M, Stampar F. Phenolic acids and flavonoids of fig fruit (*Ficus carica* L.) in northern Mediterranean region. Food Chemistry. 2008; 106:153-157.
11. Saklani S, Chandra S. Antimicrobial activity, nutritional profile and quantitative study of different fractions of *Ficus palmata*. International Research Journal of Plant Science. 2011a; 2(11):332-337.
12. Joshi Y, Joshi AK, Prasad N, Juyal D. A review on *Ficus palmate* (Wild Himalayan Fig). The Journal of Phytopharmacology. 2014; 3(5):374-377.
13. Saklani S, ChandraS. In vitro antimicrobial activity, nutritional profile and phytochemical screening of wild edible fruit of Garhwal Himalaya (*Ficus auriculata*). International Journal of Pharm Tech Research. 2012; 12(2).
14. Saklani S, Chandra S, Mishra AP. Evaluation of antioxidant activity, quantitative estimation of phenols, anthocynins and flavonoids of wild edible fruits of Garhwal Himalaya. Journal of Pharmacy Research. 2011b; 4:4083-4086.
15. AOAC. Official methods of analyses. Washington, DC: Association of Official Analytical Chemist, 1990.
16. AOAC. Official methods of analyses. Washington, DC: Association of Official Analytical Chemist, 1997.
17. Ranganna S. Handbook of Analysis and Quality Control for Fruit and Vegetable Products. 2ndedn, Tata McGraw Hill Publ. Co, New Delhi, 1986.
18. Al-Farsi M, Alasalvar C, Morris A, Barron M, Shahidi F. Comparison of anti-oxidant activity, anthocyanins, carotenoids, and phenolics of three

- native fresh and sun-dried date (*Phoenix dactylifera* L.). Journal of Agricultural and Food Chemistry. 2005; 53:7592-7599.
19. Thimmaiah SK. Standard methods of biochemical analysis (1st edn). Kalyani Publishers, New Delhi, 1999.
  20. Brand-Williams W, Cuvelier ME, Berset C. Use of a radical method to evaluate antioxidant activity. *Leben. Wiss .U. Tech.* 1995; 28:25-30.
  21. Cheftel JC, Cheftel H. Introduction à la biochimie et la technologie des aliments. Paris: Tec-Doc Lavoisier. 1976; pp: 29-171.

**HOW TO CITE THIS ARTICLE**

Kumari K, Sharma S, Joshi VK, Sharma S. Adding value to wild Himalayan fig (*Ficus palmata*): Composition, functional and sensory characteristics of jam. *J Phytopharmacol* 2018; 7(1):13-18.